

Tables and Figures for download

**National Joint Registry
for England and Wales
7th Annual Report**

Instructions:

The following lists are the same as the tables and figures lists in the annual report.

Locate the table or figure you require and then click on the JPEG or PDF button to download a copy to your desk top.

Part 1 tables

Table 1.1	List of non-returning units, 2009/10..	23	JPEG	PDF
-----------	--	----	----------------------	---------------------

Part 1 figures

Figure 1.1	NJR compliance: 2005/06 to 2009/10, based on levies from implant sales.	23	JPEG	PDF
Figure 1.2	NJR consent: annual analysis of total records received and those received with patient consent, 2005/06 to 2009/10.	24	JPEG	PDF
Figure 1.3	NJR linkability: analysis of total records received and those for which NHS numbers have been traced, 2005/06 to 2009/10.	25	JPEG	PDF
Figure 1.4	Total hip and knee joint replacement procedures entered into the NJR, 2005/06 to 2009/10, recorded by the country in which the procedure took place.	27	JPEG	PDF
Figure 1.5	Hip and knee joint replacement procedures entered into the NJR, 2005/06 to 2009/10, recorded by procedure type.	28	JPEG	PDF
Figure 1.6	Proportion of reported procedures by type of provider, 2005/06 to 2009/10.	29	JPEG	PDF
Figure 1.7	Proportion of hip replacement procedures by type of provider, 2005/06 to 2009/10.	30	JPEG	PDF
Figure 1.8	Proportion of knee replacement procedures by type of provider, 2005/06 to 2009/10.	31	JPEG	PDF
Figure 1.9	NJR Clinician Feedback PTIR hips report.	34	JPEG	PDF

Part 2 tables

Table 2.1	Total number of hospitals and treatment centres in England and Wales able to participate in the NJR and the proportion actually participating in 2009.	50	JPEG	PDF
Table 2.2	Number of participating hospitals, according to number of procedures performed during 2009.	50	JPEG	PDF
Table 2.3	Patient characteristics and procedure details, according to type of provider for hip procedures in 2009.	54	JPEG	PDF
Table 2.4	Patient characteristics for primary hip replacement procedures in 2009, according to procedure type.	57	JPEG	PDF
Table 2.5	Age and gender for primary hip replacement patients in 2009.	58	JPEG	PDF
Table 2.6	Characteristics of surgical practice for primary hip replacement procedures in 2009, according to procedure type.	62	JPEG	PDF
Table 2.7	Thromboprophylaxis regime for primary hip replacement patients, prescribed at time of operation.	64	JPEG	PDF
Table 2.8	Reported untoward intra-operative events for primary hip replacement patients in 2009, according to procedure type.	65	JPEG	PDF
Table 2.9	Patient characteristics for hip revision procedures in 2009, according to procedure type.	71	JPEG	PDF
Table 2.10	Indication for surgery for hip revision procedures, 2006 to 2009.	72	JPEG	PDF
Table 2.11	Components removed during hip revision procedures in 2009.	72	JPEG	PDF
Table 2.12	Components used during single stage hip revision procedures in 2009.	73	JPEG	PDF
Table 2.13	Patient characteristics and procedure details, according to type of provider for knee procedures in 2009.	75	JPEG	PDF

Table 2.14	Patient characteristics for primary knee replacement procedures in 2009, according to procedure type.	77		
Table 2.15	Age and gender for primary knee replacement patients in 2009.	80		
Table 2.16	Characteristics of surgical practice for primary knee replacement procedures in 2009, according to procedure type.	84	JPEG	PDF
Table 2.17	Thromboprophylaxis regime for primary knee replacement patients, prescribed at time of operation.	86	JPEG	PDF
Table 2.18	Reported untoward intra-operative events for primary knee replacement patients in 2009, according to procedure type.	87	JPEG	PDF
Table 2.19	Patient characteristics for knee revision procedures in 2009, according to procedure type.	90	JPEG	PDF

Part 2 figures

Figure 2.1	Percentage of participating hospitals by number of procedures per annum, 2004 - 2009.	51	JPEG	PDF
Figure 2.2	Primary hip procedures by type of provider.	55	JPEG	PDF
Figure 2.3	Type of primary hip replacement procedures undertaken between 2005 and 2009.	56	JPEG	PDF
Figure 2.4(a)	Age and gender for primary hip replacement patients in 2009.	59	JPEG	PDF
Figure 2.4(b)	Age for primary hip replacement patients between 2003 and 2009.	60	JPEG	PDF
Figure 2.5	ASA grades for primary hip replacement patients between 2003 and 2009.	60	JPEG	PDF
Figure 2.6(a)	BMI for primary hip replacement patients between 2004 and 2009	61	JPEG	PDF
Figure 2.6(b)	BMI groups for primary hip replacement patients between 2004 and 2009	61	JPEG	PDF
Figure 2.7	Bone cement types for primary hip replacement procedures undertaken between 2003 and 2009.	63	JPEG	PDF
Figure 2.8	Top five cemented hip stem brands, usage trends 2003 to 2009.	66	JPEG	PDF
Figure 2.9	Top five cemented hip cup brands, usage trends 2003 to 2009.	67	JPEG	PDF
Figure 2.10	Top five cementless hip stem brands, usage trends 2003 to 2009.	67	JPEG	PDF
Figure 2.11	Top five cementless hip cup brands, usage trends 2003 to 2009.	68	JPEG	PDF
Figure 2.12	Top eight resurfacing head brands, usage trends 2003 to 2009.	69	JPEG	PDF
Figure 2.13	Femoral head size trends, 2003 to 2009.	70	JPEG	PDF
Figure 2.14	Primary knee procedures by type of provider.	76	JPEG	PDF
Figure 2.15(a)	Type of primary knee replacement procedure undertaken between 2005 and 2009.	78	JPEG	PDF
Figure 2.15(b)	Implant constraint for bicondylar primary knee replacement procedures undertaken between 2005 and 2009.	79	JPEG	PDF
Figure 2.16	Age and gender for primary knee replacement patients in 2009.	81	JPEG	PDF
Figure 2.17	ASA grades for primary knee replacement patients between 2003 and 2009.	82	JPEG	PDF
Figure 2.18(a)	BMI for primary knee replacement patients between 2004 and 2009.	82	JPEG	PDF
Figure 2.18(b)	BMI groups for primary knee replacement patients between 2004 and 2009.	83	JPEG	PDF
Figure 2.19	Bone cement types for primary knee replacement procedures undertaken between 2003 and 2009.	85	JPEG	PDF
Figure 2.20	Top five total condylar knee brands, usage trends 2003 to 2009.	87	JPEG	PDF
Figure 2.21	Top five unicondylar knee brands, usage trends 2003 to 2009.	88	JPEG	PDF
Figure 2.22	Top five patello-femoral knee brands, usage trends 2003 to 2009.	88	JPEG	PDF
Figure 2.23	Top five fixed hinged knee brands, usage trends 2003 to 2009.	89	JPEG	PDF

Part 3 tables

Table 3.1	Revision rates by prosthesis type at one, three and five years for primary hip replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	101	JPEG	PDF
Table 3.2	Revision at three and five years for primary hip replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	102	JPEG	PDF
Table 3.3	Revision rates according to stem brands for primary hip replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	105	JPEG	PDF
Table 3.4	Revision rates according to cup brands for primary hip replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	106	JPEG	PDF
Table 3.5	Revision rates according to resurfacing cup brands for primary hip resurfacing procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	107	JPEG	PDF
Table 3.6	Mortality within 90 days for patients who received a primary hip replacement between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	108	JPEG	PDF
Table 3.7	Mortality within five years for patients who received a primary hip replacement between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	109	JPEG	PDF
Table 3.8	Length of hospital stay for patients who received a primary hip replacement between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	110	JPEG	PDF
Table 3.9	Revision rates by prosthesis type at one, three and five years for primary knee replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	112	JPEG	PDF
Table 3.10	Revision at three and five years for primary knee replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	114	JPEG	PDF
Table 3.11	Revision rates at three and five years according to the most frequently used brands for knee replacement procedures, undertaken between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	117	JPEG	PDF
Table 3.12	Mortality within 90 days for patients who received a primary knee replacement between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	118	JPEG	PDF
Table 3.13	Mortality within five years for patients who received a primary knee replacement between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	119	JPEG	PDF
Table 3.14	Length of hospital stay for patients who received a primary knee replacement between 1 st April 2003 and 31 st December 2009, which were linked to a HES/PEDW episode.	120	JPEG	PDF

Part 3 figures

Figure 3.1	Flowchart illustrating linkage of NJR procedures to HES/PEDW records.	97	JPEG	PDF
Figure 3.2	Implant survival of NJR-HES/PEDW linked primary hip replacements, by prosthesis type.	101	JPEG	PDF
Figure 3.3	Implant survival of NJR-HES/PEDW linked primary hip replacements, by prosthesis type, gender and age.	103	JPEG	PDF
Figure 3.4	Implant survival of NJR-HES/PEDW linked primary knee replacements, by prosthesis type.	112	JPEG	PDF
Figure 3.5	Implant survival of NJR-HES/PEDW linked primary knee replacements, by gender and prosthesis type.	115	JPEG	PDF
Figure 3.6	Implant survival of NJR-HES/PEDW linked primary knee replacements, by age and prosthesis type.	116	JPEG	PDF